

Modeling of the catalytic of benzyland butyl alcohols etherification

A.G. Vovdenko¹, M.K. Vovdenko¹, K.F. Koledina^{1,2}, A.R. Bayguzina¹, R.I. Khusnutdinov¹

¹Institute of Petrochemistry and Catalysis, Ufa Federal Research Centre of Russian Academy of Sciences, Prospect Oktyabrya 141, Ufa, Russia, 450075

²Ufa State Petroleum Technological University, Cosmonavtov str. 1, Ufa, Russia, 450062

Abstract. On the basis of experimental data obtained at different temperatures and different initial components concentrations, mathematical model of catalytic reaction of benzylbutyl ether synthesis was developed. Value of kinetic parameters, such as the rate constants and activation energies of reaction stages are determined. Thus, the developed mathematical model within the limits of the permissible error allows to adequately describe the experimental data.

1. Introduction

Benzyl butyl ether is a valuable aromatic substance and is widely used for flavoring products in the perfumery, cosmetic and food industries. Benzyl butyl ether usage use as a food flavoring (ice cream, ice, beverages, desserts, baking, etc.) is permitted in many countries. Benzyl butyl ether is a large-tonnage industrial product [1, 2].

The synthesis of benzylbutyl ether was performed by reaction between benzyl alcohol and n-butanol. Copper bromide (II) (CuBr₂) was used as a catalyst [3-7] at temperature within 140-175°C at time range of 2-10 hours at different molar ratios of initial components ([CuBr₂]:[BuOH]:[n-C₄H₉OH] = 1:100:100-300).

Mathematical model of the reaction of benzylbutyl ether synthesis based on reaction scheme [8] was developed. Chosen reaction mechanism is represented below in Table 1.

Table 1. Scheme of the reaction of benzylbutyl ether synthesis.

№	Scheme
1	$\text{PhCH}_2\text{OH}(\mathbf{X}_1) + \text{CuBr}_2(\mathbf{X}_2) \rightarrow [\text{PhCH}_2]^+[\text{CuBr}_2(\text{OH})]^- (\mathbf{X}_3)$
2	$[\text{PhCH}_2]^+[\text{CuBr}_2(\text{OH})]^- (\mathbf{X}_3) + \text{BuOH}(\mathbf{X}_4) \rightarrow [\text{PhCH}_2\text{OBu}]^+ \text{H}^+ [\text{CuBr}_2(\text{OH})]^- (\mathbf{X}_5)$
3	$[\text{PhCH}_2\text{OBu}]^+ \text{H}^+ [\text{CuBr}_2(\text{OH})]^- (\mathbf{X}_5) \rightarrow \text{PhCH}_2\text{OBu}(\mathbf{X}_6) + \text{H}_2\text{O}(\mathbf{X}_7) + \text{CuBr}_2(\mathbf{X}_2)$
4	$[\text{PhCH}_2]^+[\text{CuBr}_2(\text{OH})]^- (\mathbf{X}_3) + \text{PhCH}_2\text{OH}(\mathbf{X}_1) \rightarrow [\text{PhCH}_2\text{OHCH}_2\text{Ph}]^+ [\text{CuBr}_2(\text{OH})]^- (\mathbf{X}_8)$
5	$[\text{PhCH}_2\text{OHCH}_2\text{Ph}]^+ [\text{CuBr}_2(\text{OH})]^- (\mathbf{X}_8) \rightarrow \text{PhCH}_2\text{OCH}_2\text{Ph}(\mathbf{X}_9) + \text{H}_2\text{O}(\mathbf{X}_7) + \text{CuBr}_2(\mathbf{X}_2)$
6	$\text{BuOH}(\mathbf{X}_4) + \text{CuBr}_2(\mathbf{X}_2) \rightarrow [\text{Bu}]^+[\text{CuBr}_2(\text{OH})]^- (\mathbf{X}_{10})$
7	$[\text{Bu}]^+[\text{CuBr}_2(\text{OH})]^- (\mathbf{X}_{10}) + \text{BuOH}(\mathbf{X}_4) \rightarrow [\text{BuOHBu}]^+[\text{CuBr}_2(\text{OH})]^- (\mathbf{X}_{11})$
8	$[\text{BuOHBu}]^+[\text{CuBr}_2(\text{OH})]^- (\mathbf{X}_{11}) \rightarrow \text{BuOBu}(\mathbf{X}_{12}) + \text{H}_2\text{O}(\mathbf{X}_7) + \text{CuBr}_2(\mathbf{X}_2)$
9	$[\text{Bu}]^+[\text{CuBr}_2(\text{OH})]^- (\mathbf{X}_{10}) + \text{PhCH}_2\text{OH}(\mathbf{X}_1) \rightarrow [\text{PhCH}_2\text{OBu}]^+ \text{H}^+ [\text{CuBr}_2(\text{OH})]^- (\mathbf{X}_5)$

2. Kinetic model

2.1 Mathematic model

Using reactionscheme represented above (Table 1) mathematic model was established, based on system of differential equations:

$$\frac{dc_i}{dt} = \sum_{j=1}^J v_{ij} w_j(k_j, k_{0j}, E_j, T, c_i), \quad i = 1, \dots, I \quad (1)$$

where W_j – rate of j -th elementary reaction, [mole*min], v_{ij} – stoichiometric coefficient of i -th substance in j -th reaction, j – number of elementary reactions in kinetic model ($j = \overline{1,9}$), k_j – constant of j -th elementary reaction rate, c_i – concentration of i -th substance, [mole/lit], i – number of substance, I – overall number of components in the system ($I=12$), k_{j0} – preexponential coefficient for j -th constant of elementary reaction rate, T – temperature, E_j – activation energy of j -th reaction [KJ/mole].

After obtaining k_j values at different temperatures k_{j0} and E_j were determined from Arrhenius equation. Condition of minimization of the functional (2) was used [9-15].

$$F = \sum_{i=1}^N \sum_{j=1}^M |x_{ij}^e - x_{ij}^c| \quad (2)$$

where x_{pi}^e and x_{pi}^r are the experimental and calculated values of concentrations of the components.

2.2 Kinetic model

Using mathematical model (1), (2) and reaction scheme represented (Table 1) we solved inverse and direct kinetic tasks. In order to do this genetic algorithm and Runge-Kutta of 4th order method were used which were implemented in MATLAB software environment. Minimizing the functional (2) the kinetic parameters for different temperatures were obtained (Table 2).

Table 2. The values of the kinetic parameters of the reaction of the synthesis of benzylbutyl ether.

N stage	k_j		
	$T=140^\circ\text{C}$	$T=160^\circ\text{C}$	$T=175^\circ\text{C}$
1	0.0851	0.1134	0.2268
2	1.0280	3.0560	4.7289
3	0.1013	0.8000	1.0773
4	0.9612	2.2108	5.1905
5	0.2260	0.7000	0.8500
6	0.0003	0.0006	0.0009
7	1.1699	2.3399	4.6797
8	0.0001	0.0002	0.0004
9	0.0750	0.1500	0.3400




Figure 1. Graphs of correspondence of experimental data (points) and calculated values (lines) of changes in the concentration of the observed substrates at the ratio $[\text{CuBr}_2]:[\text{BnOH}]:[\text{n-C}_4\text{H}_9\text{OH}] = 1:100:300$, $T = 140^\circ\text{C}$.

On Figure1 – Figure 6 below are represented calculated and experimental concentrations profiles for X1, X2, X9at different temperatures and molar ratios of initial compounds.




Figure 2. Graphs of correspondence of experimental data (points) and calculated values (lines) of changes in the concentration of the observed substrates at the ratio $[CuBr_2]:[BnOH]:[n-C_4H_9OH] = 1:100:300$, $T = 160^\circ C$.




Figure 3. Graphs of correspondence of experimental data (points) and calculated values (lines) of changes in the concentration of the observed substrates at the ratio $[CuBr_2]:[BnOH]:[n-C_4H_9OH] = 1:100:300$, $T = 175^\circ C$.




Figure 4. Graphs of correspondence of experimental data (points) and calculated values (lines) of changes in the concentration of the observed substrates at the ratio $[CuBr_2]:[BnOH]:[n-C_4H_9OH] = 1:100:150$, $T = 175^\circ C$.




Figure 5. Graphs of correspondence of experimental data (points) and calculated values (lines) of changes in the concentration of the observed substrates at the ratio $[\text{CuBr}_2]:[\text{BnOH}]:[\text{n-C}_4\text{H}_9\text{OH}] = 1:100:200$, $T = 175\text{ }^\circ\text{C}$.




Figure 6. Graphs of correspondence of experimental data (points) and calculated values (lines) of changes in the concentration of the observed substrates at the ratio $[\text{CuBr}_2]:[\text{BnOH}]:[\text{n-C}_4\text{H}_9\text{OH}] = 1:100:150$, $T = 140\text{ }^\circ\text{C}$.




Figure 7. Graphs of correspondence of experimental data (points) and calculated values (lines) of changes in the concentration of the observed substrates at the ratio $[\text{CuBr}_2]:[\text{BnOH}]:[\text{n-C}_4\text{H}_9\text{OH}] = 1:100:200$, $T = 140\text{ }^\circ\text{C}$.




Figure 8. Graphs of correspondence of experimental data (points) and calculated values (lines) of changes in the concentration of the observed substrates at the ratio $[CuBr_2]:[BnOH]:[n-C_4H_9OH] = 1:100:150$, $T = 160\text{ }^\circ\text{C}$.




Figure 9. Graphs of correspondence of experimental data (points) and calculated values (lines) of changes in the concentration of the observed substrates at the ratio $[CuBr_2]:[BnOH]:[n-C_4H_9OH] = 1:100:200$, $T = 160\text{ }^\circ\text{C}$.

In Table 3 below values of activation energies represented along with values of preexponential coefficients. There also represented values of activation energies which we obtained in earlier research [7].

Table 3. Activation energies and preexponential coefficient values[8].

N stage	E_j , kcal/mol	K_{0j}	E_j , kcal/mol
1	9.93	14200.04	5.36
2	16.27	438995622.7	12.18
3	25.68	4.70666E+12	10.31
4	17.51	1676543382	13.96
5	14.35	9722953.614	21.69
6	11.60	435.72	15.03
7	14.40	47204510.15	18.46
8	14.40	3952.09	35.10
9	11.60	435.72	11.94

3. Conclusion

In this paper kinetic parameters for represented reaction scheme of benzyl butyl ether synthesis were determined. Calculations were based on experimental data which was obtained from experiments conducted at different temperatures and molar ratios of initial components. Kinetic parameters were compared with previous results [8]. It was established that developed kinetic model gives more accurate results at lower temperatures (140-160°C) and at molar ratio of initial compounds: $[CuBr_2]:[BnOH]:[n-$

C_4H_9OH] = 1:100:150-200. In order to get more accurate results at higher temperatures it is suggested to include mass exchange factors in model and make additional research in order to establish are any of elementary reactions (Table 1) of equilibrium type.

4. Acknowledgments

Reported study was funded by RSF according to the research project № 19-71-00006 (paragraph 2, 3), RFBR according to the research project № 18-07-00341.

5. References

- [1] Voytkovich, S.A. 895 fragrances for perfumes and household chemicals – Moscow: Food industry, 1994. – 594 p.
- [2] Burdock, G.A. Encyclopedia of food and color additives / G.A. Burdock – CRC Press, 1997. – 264.
- [3] Ochiai, M. Hypervalent (tert-Butylperoxy) iodanes Generate Iodine-Centered Radicals at Room Temperature in Solution: Oxidation and Deprotection of Benzyl and Allyl Ethers, and Evidence for Generation of α -Oxy Carbon Radicals / M. Ochiai, T. Ito, H. Takahashi, A. Nakanishi, M. Toyonari, T. Sueda, S. Goto, M. Shiro // *J. Am. Chem. Soc.* – 1996. – 118(33). – P. 7716-7730.
- [4] Firouzabadi, H. Facile preparation of symmetrical and unsymmetrical ethers from their corresponding alcohols catalyzed by aluminumdodecatangstophosphate ($AlPW_{12}O_{40}$), as a versatile and a highly water tolerant Lewis acid / H. Firouzabadi, N. Iranpoor, A. Jafari // *J. Mol. Catal. Chem.* – 2005. – Vol.227(1-2). – P. 97-100.
- [5] Bayguzina, A.R. Synthesis of Benzyl Alkyl Ethers by Intermolecular Dehydration of Benzyl Alcohol with Aliphatic Alcohols under the Effect of Copper Containing Catalysts / A.R. Bayguzina, L.I. Gimaletdinova, R.I. Khusnutdinov // *Russ. J. Org. Chem.* – 2018. – Vol. 54(8) – P. 1148-1155.
- [6] Liu, Y. Rhenium(I)-Catalyzed Formation of a Carbon–Oxygen Bond: An Efficient Transition Metal Catalytic System for Etherification of Benzyl Alcohols / Y. Liu, R. Hua R, H.-B. Sun, X. Qiu // *Organometallics.* – 2005. – Vol. 227(1-2). – P. 2819-2821.
- [7] Koledina, K.F. Mathematical modeling of the synthesis reaction benzyl butyl ether / K.F. Koledina, S.N. Koledin, A.G. Vovdenko, A.R. Bayguzina, R.I. Khusnutdinov // *Journal of Physics: Conference Series.* – 2019. – Vol. 1368. DOI: 10.1088/1742-6596/1368/4/042019.
- [8] Nurislamova, L. Kinetic model of the catalytic hydroalumination of olefins with organoaluminum compounds / L. Nurislamova, I. Gubaydullin, K. Koledina, R. Safin // *Reaction Kinetics, Mechanisms and Catalysis.* – 2016 – Vol. 117(1). – P. 1-14.
- [9] Koledina, K.F. Kinetics and the mechanism of the catalytic reaction of alcohols with dimethyl carbonate / K.F. Koledina, S.N. Koledin, N.A. Schadneva, I.M. Gubaidullin // *Journal of Physical Chemistry.* – 2017. – Vol. 91(3). – P. 422-428.
- [10] Koledina, K.F. Kinetic model of the catalytic reaction of dimethylcarbonate with alcohols in the presence of $Co_2(CO)_8$ and $W(CO)_6$ / K.F. Koledina, S.N. Koledin, N.A. Schadneva, Y.Yu. Mayakova, I.M. Gubaydullin // *Reaction Kinetics, Mechanisms and Catalysis.* – 2017. – Vol. 121(2). – P. 425-428.
- [11] Samarsky, A. Numerical methods / A.A. Samarsky, A.V. Gulin – Moscow "Science", 1989. – 432 p.
- [12] Gubaydullin, I.M. Informational and analytical system of inverse kinetic tasks / I.M. Gubaydullin, L.V. Sayfullina, M.R. Enikeev – Ufa: Publishing office of Bashk. University. – 2003. – 89 p.
- [13] Batishchev, D.I. Application of genetic algorithms for solving problems discrete optimization / D.I. Batishchev, E.A. Neymark, N.V. Starostin – Nizhny Novgorod, 2007. – P. 85.
- [14] Valov, A.V. Numerical methods for solving equations for engineers: tutorial material / A.V. Valov – SUSU Publishing Center, 2012. – P. 110.
- [15] Karpenko, A.P. Main essences of population algorithms for the global optimization problem // *Information and mathematical technologies in science and management.* – 2016. – Vol. 2. – P. 8-17.